



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

From (2),

$$\sin \theta = \frac{G \cos \varphi}{g} = \frac{32.05209 \times .731}{32} = .73219.$$


$$\therefore \theta = 47^\circ 4' 15''.$$

$$\therefore \text{Difference in apparent weight} = 481.0864 \cos(47^\circ 4' 15'') = 327.6583 \text{ lbs.}$$

II Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

This problem depends on the velocities in space of the Express east and the Express west.

In the figure, let P be the point of the train on the 47th parallel, O the center of the earth, AP the normal at P , φ the angle PO makes with the equatorial diameter $B'O$, θ the angle the normal makes with the same line, (θ is the latitude of P).



Let $OB = \rho$, $OP = r$, 300 tons = W , f = centrifugal force in the direction AP (vertical direction), g = gravity on 47th parallel, G = gravity at equator. $a = 6377377$ meters = 20923536 feet = equatorial radius, e = ellipticity of the earth.

$$\rho = r \cos \varphi = a \cos \theta / \sqrt{1 - e^2 \sin^2 \theta}.$$

$$\text{Now } \theta = 47^\circ, e^2 = .006920928. \therefore \rho = 4357445.45 \text{ meters.}$$

$$\text{One day} = 86400 \text{ seconds.}$$

$$\therefore 2\pi\rho/86400 = 316.8831 \text{ meters} = 1039.37 \text{ feet per second, the velocity of } P \text{ due to the earth's rotation.}$$

$$60 \text{ miles an hour} = 88 \text{ feet per second.}$$

$$1039.37 - 88 = 951.37, \text{ the train's velocity in space going west.}$$

$$1039.37 + 88 = 1127.37, \text{ the train's velocity in space going east.}$$

$$f = Wv^2/g\rho = Wv^2/gr \cos \varphi.$$

$$\therefore F = f \cos \theta = Wv^2 \sqrt{1 - e^2 \sin^2 \theta} / ag.$$

$$\text{Now } g = G(1 + \frac{1}{2}e^2 \sin^2 \theta). * \quad G = 32.2015235 \text{ feet.}$$

$$\therefore, g = 32.23130991 \text{ feet per second. } \therefore F = .000000444v^2 \text{ tons.}$$

$$\therefore F = .4018399 \text{ tons going west. } F = .5643076 \text{ tons going east.}$$

$$\therefore \text{Difference} = .1624657 \text{ tons} = 324.9354 \text{ pounds.}$$

*This expression for the true value of gravity in latitude θ is new to me. If any reader of the MONTHLY can tell me where to find it used previously, and by whom, I would be greatly pleased to know. I believe it to be new and unused before.

AVERAGE AND PROBABILITY.

76. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy, Irving College, Mechanicsburg, Pa.

In a given ellipse, the extremities of a focal chord are joined with the center. Find the average area of the angle thus formed.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.; GEORGE R. DEAN, Professor of Mathematics, University of Missouri School of Mines and Metallurgy, Rolla, Mo.; WALTER H. DRANE, Graduate Student, Harvard University, Cambridge, Mass.; J. SCHEFFER, A. M., Hagerstown, Md.; and L. C. WALKER, Instructor in Mathematics, Leland Stanford, Jr., University, Palo Alto, Cal.

Let θ = the angle the focal chord makes with the major axis.

The length, l , of this focal chord

$$= \frac{a(1-e)^2}{1+e\cos\theta} + \frac{a(1-e^2)}{1+e\cos\theta} = \frac{2a(1-e^2)}{1-e^2\cos^2\theta}.$$

The perpendicular distance from the center to this chord $= a\sin\theta$.

\therefore Area of triangle =

$$A = \frac{a^2 e(1-e^2)\sin\theta}{1-e^2\cos^2\theta}.$$

I. When the chords are drawn at equal angular intervals,

$$\Delta = \frac{\int_0^{\frac{1}{2}\pi} A d\theta}{\int_0^{\frac{1}{2}\pi} d\theta} = \frac{2a^2 e(1-e^2)}{\pi} \int_0^{\frac{1}{2}\pi} \frac{\sin\theta d\theta}{1-e^2\sin^2\theta} = \frac{a^2(1-e^2)}{\pi} \log\left(\frac{1+e}{1-e}\right).$$

DEAN, DRANE, SCHEFFER, WALKER.

II. When the abscissas of the extremity of the chord are drawn at equal intervals,

$$\begin{aligned} \Delta &= \frac{\int A dx}{\int dx} = \frac{\int_0^{\frac{1}{2}\pi} \frac{A \sin\theta d\theta}{(1-e\cos\theta)^2}}{\int_0^{\frac{1}{2}\pi} \frac{\sin\theta d\theta}{(1-e\cos\theta)^2}} = a^2 e(1-e)^2(1+e) \int_0^{\frac{1}{2}\pi} \frac{\sin^2\theta d\theta}{(1-e\cos\theta)^3(1+e\cos\theta)} \\ &= -\frac{1}{2}a^2(1-e)^2(1+e) + \frac{1}{2}a^2(1-e)^2(1+e) \int_0^{\frac{1}{2}\pi} \frac{(e+\cos\theta)d\theta}{(1-e^2\cos^2\theta)^2} \\ &= \frac{a^2(1-e)}{8b} \left(\pi a^3 e^3 - 4b(1-e^2) + 20a^2 \sqrt{1-e^2} + \frac{2a^2}{e} \sin^{-1}e \right). \end{aligned}$$

III. When the chord varies with the arc,

$$\Delta = \frac{\int A ds}{\int ds}.$$

ZERR.